ISSN (print) 0093-4666 © 2015. Mycotaxon, Ltd. ISSN (online) 2154-8889

MYCOTAXON

http://dx.doi.org/10.5248/130.1135

Volume 130, pp. 1135–1139 *October–December 2015*

Synchaetomella aquatica sp. nov. from submerged leaves from Brazil

Patrícia Oliveira Fiuza[1], Luis Fernando Pascholati Gusmão[1]* & Rafael F. Castañeda-Ruiz[2]

[1]*Departamento de Ciências Biológicas, Laboratório de Micologia, Universidade Estadual de Feira de Santana, Avenida Transnordestina s/n, Novo Horizonte, 44036-900, Feira de Santana, Brazil*

[2]*Instituto de Investigaciones Fundamentales en Agricultura Tropical 'Alejandro de Humboldt' (INIFAT), Académico Titular de la 'Academia de Ciencias de Cuba' Calle 1 Esq. 2, Santiago de Las Vegas, C. Habana, Cuba, C.P.*

*Correspondence to: *lgusmao@uefs.br*

Abstract — *Synchaetomella aquatica,* a new microfungus collected on the submerged decaying leaves of *Calophyllum brasiliense*, is described and illustrated. It is distinguished by conidiogenous cells that are synnematal, determinate conidiomata, monophialidic, discrete, and terminal and solitary falcate unicellular hyaline conidia.

Key words — asexual fungi, systematics, freshwater fungi

Introduction

During the studies of saprobic conidial fungi colonizing submerged leaves of *Calophyllum brasiliense* in the Brazilian semiarid Caatinga biome, an interesting fungus belonging to the genus *Synchaetomella* was collected. Critical morphological observation of the specimen revealed it to be an undescribed species, which we describe here as *S. aquatica*.

Materials & methods

During several expeditions between November 2013 to May 2014 through "Serra da Tromba" in the semiarid region northeastern Brazil, samples of submerged leaves of *Calophyllum brasiliense* were stored in litter bags, taken to the laboratory, and processed according to Castañeda Ruiz (2005). Mounts were prepared in PVL (polyvinyl alcohol and lactic acid) and measurements were made at a magnification of ×1000. Micrographs were obtained with an Olympus microscope (model BX51) with bright field and Nomarski interference optics. The type specimen of the new species is deposited in the Herbarium of "Universidade Estadual de Feira de Santana" (HUEFS).

Taxonomy

Synchaetomella aquatica Fiuza, Gusmão & R.F. Castañeda, **sp. nov.** Figs 1, 2

MycoBank MB812900

Differs from all other *Synchaetomella* spp. by its exclusively terminal conidiogenous cells.

Type: Brazil, Bahia, Piatã, Chapada Diamantina, Serra da Tromba, 13°07′S 41°50′W, on decaying submerged leaves of *Calophyllum brasiliense* Cambess. (*Calophyllaceae*), 11.XI.2013, coll. P.O. Fiuza (**Holotype**, HUEFS 209015).

Etymology: Latin, *aquatica*, referred to its growing in water.

On the natural substratum mycelium partly immersed and partly superficial. Hyphae septate, branched, smooth, pale brown, 2–3 µm diam. Conidiomata synnematal, determinate, scattered, erect, cylindrical, dark brown, 60–150 µm tall, 25–30 µm wide. Stipe composed of parallel hyphal filaments (conidiophores), multi-septate, straight, cylindrical, closely packed at the base and slightly loosely and branched towards the apex, brown below, brown or pale brown above, smooth, 1.5–3 µm diam. Conidiogenous cells monophialidic, discrete, terminal, determinate, brown to pale brown, 15–45 × 1.5–2.5 µm with a narrow, slightly obscure, sub-cupulate collarette with a distinct, very narrow periclinal channel, 1–2.5 µm deep. Conidia solitary, acrogenous, falcate, unicellular, 8–18 × 1.5–2 µm, smooth, hyaline, accumulating in a white mucilaginous mass.

Note: Decock et al. (2005) introduced the synnematous genus *Synchaetomella* Decock & Seifert, typified by *S. lunatospora* Decock et al. and characterized by phialidic, terminal and intercalary conidiogenous cells and falcate, 0–1-septate, hyaline conidia. The second species, *S. acerina* (Seifert & Louis-Seize 2012), also has intercalary and terminal conidiogenous cells, but the conidia are unicellular, allantoid to ellipsoidal. *Synchaetomella aquatica* is clearly differentiated by the terminal subulate conidiogenous cells.

Acknowledgments

The authors express their sincere gratitude to Dr. Xiu-Guo Zhang and Dr. De-Wei Li for their critical review of the manuscript. The authors are grateful to Dr. Keith Seifert for his commentaries and opinion. POF thanks the 'Coordenação de Aperfeiçoamento de Pessoal de Nível Superior" (CAPES) and the 'Programa de Pós-graduação em Botânica PPGBot/UEFS'. The authors are grateful to the "Programa de Pesquisa em Biodiversidade" – (PPBio Semi-arid/MCTI/CNPq) for financial support. The authors thank the support provided by "Programa Ciência sem Fronteiras". LFPG is grateful

Fig. 1. *Synchaetomella aquatica* (ex holotype HUEFS 209015). A, B. conidia; C, D. conidiogenous cells and conidia. Scale bars: A, C, D = 10 µm, B = 20 µm.





Fig 2. *Synchaetomella aquatica* (ex holotype HUEFS 209015).
A. conidia; B. conidiogenous cells and conidia; C. synnema.
Scale bars: A, B = 10 µm; C = 25 µm.

to CNPq for grant (Proc. 303062/2014-2) and RFCR is grateful to Cuban Ministry of Agriculture and "Programa de Salud Animal y Vegetal", project P131LH003033 for facilities. We acknowledge the facilities provided by Dr. P.M. Kirk and Drs. V. Robert and A. Decock through the Index Fungorum and MycoBank websites. Dr. Lorelei Norvell's editorial review and Dr. Shaun Pennycook's nomenclature review are greatly appreciated.

Literature cited

Castañeda-Ruiz RF. 2005. Metodología en el estudio de los hongos anamorfos. 182–183, in: Anais do V Congresso Latino Americano de Micología. Brasilia.

Decock C, Delgado RG, Seifert K. 2005. Taxonomy and phylogeny of *Synchaetomella lunatospora*, a new genus and species of synnematous fungi from Southeast Asia. Antonie van Leeuwenhoek 88: 231–240. http://dx.doi.org/10.1007/s10482-005-7210-7

Seifert KA, Louis-Seize G. 2012. *Synchaetomella acerina* Seifert, sp. nov. (Fungal Planet 150). Persoonia 29: 192–193. http://dx.doi.org/10.3767/003158512X661589